

MICROWAVE SPECTROSCOPIC AND THEORETICAL EXAMINATION OF ALLYL ISOTHIOCYANATE CONFORMATIONAL SPACE

JOSEPH STITSKY, WENHAO SUN, WESLEY G. D. P. SILVA, JENNIFER VAN WIJNGAARDEN,
Department of Chemistry, University of Manitoba, Winnipeg, MB, Canada.

A spectroscopic analysis of allyl isothiocyanate ($\text{CH}_2 = \text{CHCH}_2\text{NCS}$) was conducted via chirped-pulse and Balle-Flygare Fourier Transform microwave (FTMW) spectroscopy in the 7-25 GHz range. Rotational transitions associated with the different conformers were assigned based on information gathered from quantum-chemical calculations computed at B3LYP-D3(BJ) and MP2 levels of theory using the Dunning cc-pVTZ basis set. The results showed the existence of two conformers, the gauche conformer, which was reported in previous work^a as well as the first reported existence of a second conformer which corresponds to the global minimum. Spectroscopic assignments included transitions due to both parent conformers, ³⁴S, ¹³C and ¹⁵N singly substituted isotopic species as well as hyperfine splitting due to the presence of ¹⁴N.

^aMaiti, S.; Jaman, A. I.; Nandi, R. N. *J. Mol. Spectrosc.* **165** (1994), 168-172